

Chapter 6

Universality¹

A. B. J. KUIJLAARS²

Abstract

Universality of eigenvalue spacings is one of the basic characteristics of random matrices. We give the precise meaning of universality and discuss the standard universality classes (sine, Airy, Bessel) and their appearance in unitary, orthogonal, and symplectic ensembles. The Riemann-Hilbert problem for orthogonal polynomials is one possible tool to derive universality in unitary random matrix ensembles. An overview is presented of the Deift/Zhou steepest descent analysis of the Riemann-Hilbert problem in the one-cut regular case. Non-standard universality classes that arise at singular points in the spectrum are discussed at the end.

6.1 Heuristic meaning of universality

Universality of eigenvalue spacings is one of the basic characteristic features of random matrix theory. It was the main motivation of Wigner to introduce random matrix ensembles to model energy levels in quantum systems.

On a local scale the eigenvalues of random matrices show a repulsion which implies that it is highly unlikely that eigenvalues are very close to each other. Also very large gaps between neighboring eigenvalues are unlikely. This is in sharp contrast to points that are sampled independently from a given distribution. Such points exhibit Poisson statistics in contrast to what is now called

¹To appear as Chapter 6 in: Oxford Handbook of Random Matrix Theory, (G. Akemann, J. Baik, and P. Di Francesco, eds.), Oxford University Press, 2011.

² Department of Mathematics, Katholieke Universiteit Leuven, Celestijnenlaan 200B, 3001 Leuven, Belgium. The author is supported in part by FWO-Flanders projects G.0427.09 and G.0641.11, by K.U. Leuven research grant OT/08/33, by the Belgian Interuniversity Attraction Pole P06/02, and by a grant from the Ministry of Education and Science of Spain, project code MTM2005-08648-C02-01.

GUE/GOE/GSE statistics. These new random matrix theory type statistics have been observed (and sometimes proved) in many other mathematical and physical systems as well, see [Dei07d], and are therefore universal in a sense that goes beyond random matrix theory. Examples outside of random matrix theory are discussed in many chapters of this handbook. In this chapter the discussion of the universality is however restricted to random matrix theory.

The universality conjecture in random matrix theory says that the local eigenvalue statistics of many random matrix ensembles are the same, that is, they do not depend on the exact probability distribution that is put on a set of matrices, but only on some general characteristics of the ensemble. The universality of local eigenvalue statistics is a phenomenon that takes place for large random matrices. In a proper mathematical formulation it is a statement about a certain limiting behavior as the size of the matrices tends to infinity.

The characteristics that play a role in the determination of the universality classes are the following.

- Invariance properties of the ensemble, of which the prototype is invariance with respect to orthogonal, unitary, or unitary-symplectic conjugation. As discussed in Chapters 4 and 5 of this handbook, these lead to random matrix ensembles with an explicit joint eigenvalue density of the form

$$p(x_1, \dots, x_n) = \frac{1}{Z_{n,\beta}} \prod_{1 \leq i < j \leq n} |x_j - x_i|^\beta \prod_{j=1}^n e^{-V(x_j)} \quad (6.1.1)$$

where $\beta = 1, 2, 4$ corresponds to orthogonal, unitary, and symplectic ensembles, respectively. The case $V(x) = \frac{1}{2}x^2$ gives the Gaussian ensembles GOE, GUE, and GSE. The local eigenvalue repulsion increases with β and different values of β give rise to different universality results.

- An even more basic characteristic is the distinction between random matrix ensembles of real symmetric or complex Hermitian matrices. Even without the invariance assumption, the universality of local eigenvalue statistics is conjectured to hold. This has been proved recently for large classes of Wigner ensembles, i.e., random matrix ensembles with independent, identically distributed entries, see [Erd10a, Erd10b, Tao11] and the survey [Erd10c]. See Chapter 21 of this handbook for more details. Universality is also expected to exist in classes of non-Hermitian matrices, see Chapter 18.
- Another main characteristic is the nature of the point around which the local eigenvalue statistics are considered. A typical point in the spectrum is such that, possibly after appropriate scaling, the limiting mean eigenvalue density is positive. Such a point is in the bulk of the spectrum, and universality around such a point is referred to as bulk universality.

At edge points of the limiting spectrum the limiting mean eigenvalue density typically vanishes like a square root, and at such points a different type of universality is expected, which is known as (soft) edge universality. At natural edges of the spectrum, such as the point zero for ensembles of positive definite matrices, the limiting mean eigenvalue density typically blows up like the inverse of a square root, and universality at such a point is known as hard edge universality.

- At very special points in the limiting spectrum the limiting mean eigenvalue density may exhibit singular behavior. For example, the density may vanish at an isolated point in the interior of the limiting spectrum, or it may vanish to higher order than square root at soft edge points. This non-generic behavior may take place in ensembles of the form (6.1.1) with a varying potential NV

$$p(x_1, \dots, x_n) = \frac{1}{Z_{n,\beta}} \prod_{1 \leq i < j \leq n} |x_j - x_i|^\beta \prod_{j=1}^n e^{-NV(x_j)} \quad (6.1.2)$$

as $n, N \rightarrow \infty$ with $n/N \rightarrow 1$.

At such special points one expects other universality classes determined by the nature of the vanishing of the limiting mean eigenvalue density at that point, see also a discussion in Chapter 14.

In the rest of this chapter we first give the precise meaning of the notion of universality and we discuss the limiting kernels (sine, Airy and Bessel) associated with the bulk universality and the edge universality for the unitary, orthogonal, and symplectic universality classes. In Section 6.3 we discuss unitary matrix ensembles in more detail and we show that universality in these ensembles comes down to the convergence of the properly scaled eigenvalue correlation kernels. In Section 6.4 we discuss the Riemann-Hilbert method in some detail. The Riemann-Hilbert method is one of the main methods to prove universality in unitary ensembles. In the final Section 6.5 we discuss certain non-standard universality classes that arise at singular points in the limiting spectrum. We describe the limiting kernels for each of the three types of singular points, namely interior singular points, singular edge points, and exterior singular points.

Other approaches to universality are detailed in Chapter 21 for Wigner ensembles and in chapter 16 using loop equation techniques.

6.2 Precise statement of universality

For a probability density $p_n(x_1, \dots, x_n)$ on $\mathbb{R}^n$ that is symmetric (i.e., invariant under permutations of coordinates), let us define the k -point correlation

function by

$$R_{n,k}(x_1, \dots, x_k) = \frac{n!}{(n-k)!} \int \cdots \int p_n(x_1, \dots, x_n) dx_{k+1} \cdots dx_n. \quad (6.2.1)$$

Up to the factor $n!/(n-k)!$ this is the k th marginal distribution of p_n .

Fix a reference point x^* and a constant $c_n > 0$. We center the points around x^* and scale by a factor c_n , so that $(x_1, \dots, x_n)$ is mapped to

$$(c_n(x_1 - x^*), \dots, c_n(x_n - x^*)).$$

These centered and scaled points have the following rescaled k -point correlation functions

$$\frac{1}{c_n^k} R_{n,k} \left(x^* + \frac{x_1}{c_n}, x^* + \frac{x_2}{c_n}, \dots, x^* + \frac{x_k}{c_n} \right). \quad (6.2.2)$$

The universality is a property of a sequence (p_n) of symmetric probability density functions. Universality at x^* means that for a suitably chosen sequence (c_n) the rescaled k -point correlation functions (6.2.2) have a specific limit as $n \rightarrow \infty$. The precise limit determines the universality class.

As discussed in chapter 4 for determinantal point processes, the main spectral statistics such as gap probabilities and eigenvalue spacings can be expressed in terms of the k -point correlation functions. If the limits (6.2.2) exist and belong to a certain universality class, then this also leads to the universal behavior of the local eigenvalue statistics.

6.2.1 Unitary universality classes

The unitary universality classes are characterized by the fact that the limit of (6.2.2) can be expressed as a $k \times k$ determinant

$$\det [K(x_i, x_j)]_{1 \leq i, j \leq k}$$

involving a kernel $K(x, y)$, which is called the eigenvalue correlation kernel.

Bulk universality A sequence of symmetric probability density functions (p_n) then has bulk universality at a point x^* if there exists a sequence (c_n) , so that for every k , the limit of (6.2.2) is given by

$$\det [K^{\sin}(x_i, x_j)]_{1 \leq i, j \leq k}$$

where

$$K^{\sin}(x, y) = \frac{\sin \pi(x - y)}{\pi(x - y)} \quad (6.2.3)$$

is the sine kernel. Bulk universality is also known as GUE statistics.

Edge universality The (soft) edge universality holds at x^* if the limit of (6.2.2) is equal to

$$\det [K^{\text{Ai}}(x_i, x_j)]_{1 \leq i, j \leq k}$$

where

$$K^{\text{Ai}}(x, y) = \frac{\text{Ai}(x) \text{Ai}'(y) - \text{Ai}'(x) \text{Ai}(y)}{x - y} \quad (6.2.4)$$

is the Airy kernel. Here Ai denotes the Airy function, which is the solution of the Airy differential equation $y''(x) = xy(x)$ with asymptotics

$$\text{Ai}(x) = \frac{1}{2\sqrt{\pi}x^{1/4}} e^{-\frac{2}{3}x^{3/2}} \left(1 + O(x^{-3/2})\right), \quad \text{as } x \rightarrow +\infty.$$

The Airy kernel is intimately related to the Tracy-Widom distribution for the largest eigenvalue in random matrix theory [Tra94].

Hard edge universality A hard edge is a boundary for the eigenvalues that is part of the model. For example, if the random matrices are real symmetric (or complex Hermitian) positive definite then all eigenvalues are non-negative and zero is a hard edge.

The hard edge universality holds at x^* if the limit of (6.2.2) is equal to

$$\det [K^{\text{Bes}, \alpha}(x_i, x_j)]_{1 \leq i, j \leq k}$$

where

$$K^{\text{Bes}, \alpha}(x, y) = \frac{J_\alpha(\sqrt{x})\sqrt{y}J'_\alpha(\sqrt{y}) - \sqrt{x}J'_\alpha(\sqrt{x})J_\alpha(\sqrt{y})}{2(x - y)}, \quad x, y > 0, \quad (6.2.5)$$

and J_α is the usual Bessel function of order α .

The Bessel kernels depend on the parameter $\alpha > -1$ which may be interpreted as a measure of the interaction with the hard edge. The bigger α , the more repulsion from the hard edge.

6.2.2 Orthogonal and symplectic universality classes

The orthogonal and symplectic universality classes are characterized by the fact that the limit of (6.2.2) is expressed as a Pfaffian

$$\text{Pf} [K(x_i, x_j)]_{1 \leq i, j \leq k} \quad (6.2.6)$$

of a 2×2 matrix kernel

$$K(x, y) = \begin{pmatrix} K_{11}(x, y) & K_{12}(x, y) \\ K_{21}(x, y) & K_{22}(x, y) \end{pmatrix}. \quad (6.2.7)$$

Recall that the Pfaffian $\text{Pf}(A)$ of a $2k \times 2k$ skew symmetric matrix A is such that

$$\text{Pf}(A) = \sqrt{\det(A)}.$$

The matrix in (6.2.6) is a $2k \times 2k$ matrix written as a $k \times k$ block matrix with 2×2 blocks. It is skew symmetric provided that $K(y, x) = -K(x, y)^T$ where ^{T} denotes the matrix transpose.

Bulk universality A sequence of symmetric probability density functions (p_n) has orthogonal/symplectic bulk universality at a point x^* if there exists a sequence (c_n) , so that for every k , the limit of (6.2.2) is given by

$$\text{Pf} \left[K^{\sin, \beta}(x_i, x_j) \right]_{1 \leq i, j \leq k}$$

with

$$K^{\sin, \beta=1}(x, y) = \begin{pmatrix} -\frac{\partial}{\partial x} \frac{\sin \pi(x-y)}{\pi(x-y)} & \frac{\sin \pi(x-y)}{\pi(x-y)} \\ -\frac{\sin \pi(x-y)}{\pi(x-y)} & \int_0^{x-y} \frac{\sin \pi t}{\pi t} dt - \frac{1}{2} \text{sgn}(x-y) \end{pmatrix}$$

where $\text{sgn}(t) = 1, 0, -1$ depending on whether $t > 0$, $t = 0$, or $t < 0$, in the case of orthogonal (i.e., $\beta = 1$) bulk universality, and

$$K^{\sin, \beta=4}(x, y) = \begin{pmatrix} -\frac{\partial}{\partial x} \frac{\sin 2\pi(x-y)}{2\pi(x-y)} & \frac{\sin 2\pi(x-y)}{2\pi(x-y)} \\ -\frac{\sin 2\pi(x-y)}{2\pi(x-y)} & \int_0^{x-y} \frac{\sin 2\pi t}{2\pi t} dt \end{pmatrix}$$

in the case of symplectic (i.e., $\beta = 4$) bulk universality.

Edge universality The orthogonal/symplectic (soft) edge universality holds at x^* if the limit of (6.2.2) is equal to

$$\text{Pf} \left[K^{\text{Ai}, \beta}(x_i, x_j) \right]_{1 \leq i, j \leq k}$$

with

$$\begin{aligned} K_{11}^{\text{Ai}, \beta=1}(x, y) &= \frac{\partial}{\partial y} K^{\text{Ai}}(x, y) + \frac{1}{2} \text{Ai}(x) \text{Ai}(y) \\ K_{12}^{\text{Ai}, \beta=1}(x, y) &= K^{\text{Ai}}(x, y) + \frac{1}{2} \text{Ai}(x) \cdot \int_{-\infty}^y \text{Ai}(t) dt \\ K_{21}^{\text{Ai}, \beta=1}(x, y) &= -K_{12}^{\text{Ai}, \beta=1}(x, y) \\ K_{22}^{\text{Ai}, \beta=1}(x, y) &= -\int_x^\infty K^{\text{Ai}}(t, y) dt - \frac{1}{2} \int_x^y \text{Ai}(t) dt \\ &\quad + \frac{1}{2} \int_x^\infty \text{Ai}(t) dt \cdot \int_y^\infty \text{Ai}(t) dt - \frac{1}{2} \text{sgn}(x-y) \end{aligned} \tag{6.2.8}$$

in the case of orthogonal edge universality, and

$$\begin{aligned}
K_{11}^{\text{Ai}, \beta=4}(x, y) &= \frac{1}{2} \frac{\partial}{\partial y} K^{\text{Ai}}(x, y) + \frac{1}{4} \text{Ai}(x) \text{Ai}(y) \\
K_{12}^{\text{Ai}, \beta=4}(x, y) &= \frac{1}{2} K^{\text{Ai}}(x, y) - \frac{1}{4} \text{Ai}(x) \cdot \int_y^\infty \text{Ai}(t) dt \\
K_{21}^{\text{Ai}, \beta=4}(x, y) &= -K_{12}^{\text{Ai}, \beta=4}(x, y) \\
K_{22}^{\text{Ai}, \beta=4}(x, y) &= -\frac{1}{2} \int_x^\infty K^{\text{Ai}}(t, y) dt + \frac{1}{4} \int_x^\infty \text{Ai}(t) dt \cdot \int_y^\infty \text{Ai}(t) dt
\end{aligned} \tag{6.2.9}$$

in the case of symplectic edge universality. The kernel K^{Ai} that appears in (6.2.8) and (6.2.9) is the Airy kernel from (6.2.4).

Hard edge universality The orthogonal/symplectic hard edge universality is expressed in terms of 2×2 kernels with Bessel functions. See Forrester [For10] for the precise statement.

6.2.3 Determinantal and Pfaffian point processes

The universality limits have the characteristic properties of determinantal or Pfaffian point processes, see [For10, Sos00] and Chapter 11 of this handbook.

If the probability densities p_n on $\mathbb{R}^n$ also arise from determinantal point processes, then the statement of universality can be expressed as the convergence of the corresponding correlation kernels after appropriate scaling.

The probability density p_n arises from a determinantal point process, if there exist (scalar) kernels K_n so that for every n and k ,

$$R_{n,k}(x_1, \dots, x_k) = \det [K_n(x_i, x_j)]_{1 \leq i, j \leq k}.$$

In particular, one then has for the 1-point function (particle density),

$$R_{n,1}(x) = K_n(x, x)$$

and for the probability density p_n itself

$$p_n(x_1, \dots, x_n) = \frac{1}{n!} \det [K_n(x_i, x_j)]_{1 \leq i, j \leq n}.$$

In this setting the bulk universality comes down to the statement that the centered and rescaled kernels

$$\frac{1}{c_n} K_n \left(x^* + \frac{x}{c_n}, x^* + \frac{y}{c_n} \right) \tag{6.2.10}$$

tend to the sine kernel (6.2.3) as $n \rightarrow \infty$, and likewise for the edge universality.

Similarly, the probability densities p_n come from Pfaffian point processes, if there exist 2×2 matrix kernels K_n so that for every n and k ,

$$R_{n,k}(x_1, \dots, x_k) = \text{Pf} [K_n(x_i, x_j)]_{1 \leq i, j \leq k}.$$

A proof of universality in a sequence of Pfaffian point processes then comes down to the proof of a scaling limit for the matrix kernels as $n \rightarrow \infty$.

Unitary random matrix ensembles are basic examples of determinantal point processes, while orthogonal and symplectic matrix ensembles are examples of Pfaffian point processes.

For the classical ensembles that are associated with Hermite, Laguerre and Jacobi polynomials the existence and the form of the limiting kernels has been proven using the explicit formulas that are available for these classical orthogonal polynomials, see e.g. [For10, Meh04].

For non-classical ensembles, the results about universality are fairly complete for unitary ensembles, due to their connection with orthogonal polynomials. This will be discussed in more detail in the next section, and we give references there.

The first rigorous results on bulk universality for orthogonal and symplectic ensembles are due to Stojanovic [Sto00] who discussed ensembles with a quartic potential. This was extended by Deift and Gioev [Dei07a] for ensembles (6.1.1) with polynomial V . Their techniques are extended to treat edge universality in [Dei07b, Dei07c], see also the recent monograph [Dei09]. Varying weights are treated by Shcherbina in [Shc08, Shc09a].

6.3 Unitary random matrix ensembles

We explain in more detail how the universality classes arise for the eigenvalues of a unitary invariant ensemble

$$\frac{1}{\tilde{Z}_{n,N}} e^{-N \text{Tr} V(M)} dM \tag{6.3.1}$$

defined on the space of $n \times n$ Hermitian matrices, where

$$dM = \prod_{i=1}^n dM_{ii} \prod_{1 \leq i < j \leq n} d \text{Re} M_{ij} d \text{Im} M_{ij},$$

and $\tilde{Z}_{n,N}$ is a normalization constant, see also Chapter 4. The potential function V in (6.3.1) is typically a polynomial, but could be more general as well. To ensure that (6.3.1) is well-defined as a probability measure, we assume that

$$\lim_{x \rightarrow \pm\infty} \frac{V(x)}{\log(1+x^2)} = +\infty. \tag{6.3.2}$$

The factor N in (6.3.1) will typically be proportional to n . This is needed in the large n limit in order to balance the repulsion among eigenvalues due to the Vandermonde factor in (6.3.3) and the confinement of eigenvalues due to the potential V .

6.3.1 Orthogonal polynomial kernel

The joint probability density for the eigenvalues of a matrix M of the ensemble (6.3.1) has the form

$$p_{n,N}(x_1, \dots, x_n) = \frac{1}{Z_{n,N}} \prod_{1 \leq i < j \leq n} (x_i - x_j)^2 \prod_{j=1}^n e^{-NV(x_j)}. \quad (6.3.3)$$

Introduce the monic polynomials $P_{k,N}$, $P_{k,N}(x) = x^k + \dots$, that are orthogonal with respect to the weight $e^{-NV(x)}$ on $\mathbb{R}$:

$$\int_{-\infty}^{\infty} P_{k,N}(x) x^j e^{-NV(x)} dx = \begin{cases} 0, & \text{for } j = 0, \dots, k-1, \\ \gamma_{k,N}^2 > 0, & \text{for } j = k, \end{cases} \quad (6.3.4)$$

and the orthogonal polynomial kernel

$$K_{n,N}(x, y) = \sqrt{e^{-NV(x)}} \sqrt{e^{-NV(y)}} \sum_{k=0}^{n-1} \frac{P_{k,N}(x) P_{k,N}(y)}{\gamma_{k,N}^2}. \quad (6.3.5)$$

Using the formula for a Vandermonde determinant and performing elementary row operations we write (6.3.3) as

$$\begin{aligned} p_{n,N}(x_1, \dots, x_n) &= \frac{1}{Z_{n,N}} \left(\det [P_{k-1,N}(x_j)]_{1 \leq j, k \leq n} \right)^2 \prod_{j=1}^n e^{-NV(x_j)} \\ &= \frac{\prod_{k=0}^{n-1} \gamma_{k,N}^2}{Z_{n,N}} \left(\det \left[\frac{\sqrt{e^{-NV(x_j)}} P_{k-1,N}(x_j)}{\gamma_{k-1,N}} \right]_{1 \leq j, k \leq n} \right)^2. \end{aligned}$$

Evaluating the square of the determinant using the rule $(\det A)^2 = \det(AA^T)$ we obtain that (6.3.3) can be written as

$$p_{n,N}(x_1, \dots, x_n) = \frac{1}{n!} \det [K_{n,N}(x_i, x_j)]_{1 \leq i, j \leq n}, \quad (6.3.6)$$

since it can be shown that $Z_{n,N} = n! \prod_{k=0}^{n-1} \gamma_{k,N}^2$. The orthogonality condition (6.3.4) is then used to prove that the k -point correlation functions (6.2.1) are also determinants

$$R_{n,k}(x_1, \dots, x_k) = \det [K_{n,N}(x_i, x_j)]_{1 \leq i, j \leq k},$$

which shows that the eigenvalues are a determinantal point process with correlation kernel $K_{n,N}$. For the above calculation, see also Chapter 4.

The eigenvalues of orthogonal and symplectic ensembles of random matrices have joint probability density

$$\frac{1}{Z_{n,N}} \prod_{1 \leq i < j \leq n} |x_i - x_j|^\beta \prod_{j=1}^n e^{-NV(x_j)}$$

with $\beta = 1$ for orthogonal ensembles and $\beta = 4$ for symplectic ensembles. These probability densities are basic examples of Pfaffian ensembles, as follows from the calculations in e.g. [Dei09, Tra98], see also Chapter 5 of this handbook.

6.3.2 Global eigenvalue regime

As $n, N \rightarrow \infty$ such that $n/N \rightarrow 1$ the eigenvalues have a limiting distribution. A weak form of this statement is expressed by the fact that

$$\lim_{\substack{n, N \rightarrow \infty \\ n/N \rightarrow 1}} \frac{1}{n} K_{n,N}(x, x) = \rho_V(x), \quad x \in \mathbb{R} \quad (6.3.7)$$

exists. The density ρ_V describes the global or macroscopic eigenvalue regime.

The probability measure $d\mu_V(x) = \rho_V(x)dx$ minimizes the weighted energy

$$\iint \log \frac{1}{|x - y|} d\mu(x) d\mu(y) + \int V(x) d\mu(x) \quad (6.3.8)$$

among all Borel probability measures μ on $\mathbb{R}$. Heuristically, it is easy to understand why (6.3.8) is relevant. Indeed, from (6.3.3) we find after taking logarithms and dividing by $-n^2$, that the most likely distribution of eigenvalues $x_1, \dots, x_n$ for (6.3.3) is the one that minimizes

$$\frac{1}{n^2} \sum_{\substack{i, j=1 \\ i \neq j}}^n \log \frac{1}{|x_i - x_j|} + \frac{N}{n^2} \sum_{j=1}^n V(x_j).$$

This discrete minimization problem leads to the minimization of (6.3.8) in the continuum limit as $n, N \rightarrow \infty$ with $n/N \rightarrow 1$.

The minimizer of (6.3.8) is unique and has compact support. It is called the equilibrium measure in the presence of the external field V , because of its connections with logarithmic potential theory [Saf97]. The proof of (6.3.7) with $\rho_V(x)dx$ minimizing (6.3.8) is in [Dei99a, Joh98]. See also the remark at the end of subsection 6.4.8 below. It follows as well from the more general large deviation principle that is associated with the weighted energy (6.3.8), see [Ben97] and also [And10, §2.6]. See also Chapter 14 of this handbook.

If V is real analytic on $\mathbb{R}$ then μ_V is supported on a finite union of intervals, say

$$\text{supp}(\mu_V) = \bigcup_{j=1}^m [a_j, b_j],$$

and the density ρ_V takes the form

$$\rho_V(x) = \frac{1}{\pi} h_j(x) \sqrt{(b_j - x)(x - a_j)}, \quad x \in [a_j, b_j], \quad j = 1, \dots, m \quad (6.3.9)$$

where h_j is real analytic and non-negative on $[a_j, b_j]$, see [Dei98]. Another useful representation is that

$$\rho_V(x) = \frac{1}{\pi} \sqrt{q_V^-(x)}, \quad x \in \mathbb{R}, \quad (6.3.10)$$

where q_V^- is the negative part of

$$q_V(x) = \left(\frac{V'(x)}{2} \right)^2 - \int \frac{V'(x) - V'(s)}{x - s} d\mu_V(s). \quad (6.3.11)$$

If V is a polynomial then q_V is a polynomial of degree $2(\deg V - 1)$. In that case the number of intervals in the support is bounded by $\frac{1}{2} \deg V$, see again [Dei98]. See also a discussion in Chapters 14 and 16.

6.3.3 Local eigenvalue regime

In the context of Hermitian matrix models, the universality may be stated as the fact that the global eigenvalue regime determines the local eigenvalue regime. The universality results take on a different form depending on the nature of the reference point x^* .

Bulk universality A regular point in the bulk is an interior point x^* of $\text{supp}(\mu_V)$ such that $\rho_V(x^*) > 0$. At a regular point in the bulk one has

$$\lim_{n \rightarrow \infty} \frac{1}{cn} K_{n,N} \left(x^* + \frac{x}{cn}, x^* + \frac{y}{cn} \right) = K^{\sin}(x, y) \quad (6.3.12)$$

where $c = \rho_V(x^*)$ and $K^{\sin}$ is the sine kernel (6.2.3).

Convincing heuristic arguments for the universality of the sine kernel were given by Brézin and Zee [Bre93] by the method of orthogonal polynomials. Supersymmetry arguments were used in [Hac95]. Rigorous results for bulk universality (6.3.12) beyond the classical ensembles were first given by Pastur and Shcherbina [Pas97], and later by Bleher and Its [Ble99] and by Deift et al. [Dei99b, Dei99c] for real analytic V . In these papers the Riemann-Hilbert techniques were introduced in the study of orthogonal polynomials, which we

will review in section 6.4 below Conditions on V in (6.1.2) were further weakened in [McL08, Pas08].

In more recent work of Lubinsky [Lub08a, Lub09a, Lub09b] and Levin and Lubinsky [Lev08, Lev09] it was shown that bulk universality holds under extremely weak conditions. One of the results is that in an ensemble (6.1.1) restricted to a compact interval, bulk universality holds at each interior point where V is continuous.

Soft edge universality An edge point $x^* \in \{a_1, b_1, \dots, a_m, b_m\}$ is a regular edge point of $\text{supp}(\mu_V)$ if $h_j(x^*) > 0$ in (6.3.9). In that case the density ρ_V vanishes as a square root at x^* . The scaling limit is then the Airy kernel (6.2.4). If $x^* = b_j$ is a right-edge point then for a certain constant $c > 0$,

$$\lim_{n \rightarrow \infty} \frac{1}{(cn)^{2/3}} K_{n,N} \left(x^* + \frac{x}{(cn)^{2/3}}, x^* + \frac{y}{(cn)^{2/3}} \right) = K^{\text{Ai}}(x, y) \quad (6.3.13)$$

while if $x^* = a_j$ is a left-edge point, we find the same limit after a change of sign

$$\lim_{n \rightarrow \infty} \frac{1}{(cn)^{2/3}} K_{n,N} \left(x^* - \frac{x}{(cn)^{2/3}}, x^* - \frac{y}{(cn)^{2/3}} \right) = K^{\text{Ai}}(x, y). \quad (6.3.14)$$

Thus the regular edge points belong to the Airy universality class.

The Airy kernel was implicitly derived in [Bow91]. For classical ensembles the soft edge universality (6.3.13) is derived in [For93, Nag93], see also [For10]. For quartic and sextic potentials V , it is derived in [Kan97] by the so-called Shohat method. For real analytic potentials it is implicit in [Dei99c] and made explicit in [Dei07b].

Hard edge universality The Bessel universality classes arise for eigenvalues of positive definite matrices. Let $\alpha > -1$ be a parameter and consider

$$\frac{1}{\tilde{Z}_{n,N}} (\det M)^\alpha e^{-N \text{Tr } V(M)} dM$$

as a probability measure on the space of $n \times n$ Hermitian positive definite matrices. For the case $V(x) = x$ this is the Laguerre Unitary Ensemble, which is also known as a complex Wishart ensemble, see Chapter 4.

Assuming that α remains fixed, the global eigenvalue regime does not depend on α . The eigenvalue density ρ_V lives on $[0, \infty)$, and for $x \in [0, \infty)$ it continues to have the representation (6.3.10) but now q_V is modified to

$$q_V(x) = \left(\frac{V'(x)}{2} \right)^2 - \int \frac{V'(x) - V'(s)}{x - s} d\mu_V(s) - \frac{1}{x} \int V'(s) d\mu_V(s). \quad (6.3.15)$$

If $\int V'(s)d\mu_V(s) > 0$ then 0 is in the support of μ_V and ρ_V has a square-root singularity at $x = 0$.

The effect of the parameter α is noticeable in the local eigenvalue regime near $x^* = 0$. If $\int V'(s)d\mu_V(s) > 0$, then the limiting kernel is the Bessel kernel (6.2.5) of order α . For $x, y > 0$ and for an appropriate constant $c > 0$, we have

$$\lim_{n \rightarrow \infty} \frac{1}{(cn)^2} K_{n,N} \left(\frac{x}{(cn)^2}, \frac{y}{(cn)^2} \right) = K^{\text{Bes}, \alpha}(x, y). \quad (6.3.16)$$

The hard edge universality (6.3.16) was proven in [Kui02, Lub08b].

Spectral singularity A related class of Bessel kernels

$$\widehat{K}^{\text{Bes}, \alpha}(x, y) = \pi \sqrt{x} \sqrt{y} \frac{J_{\alpha+\frac{1}{2}}(\pi x) J_{\alpha-\frac{1}{2}}(\pi y) - J_{\alpha-\frac{1}{2}}(\pi x) J_{\alpha+\frac{1}{2}}(\pi y)}{2(x-y)} \quad (6.3.17)$$

appears as scaling limits in Hermitian matrix models of the form

$$\frac{1}{\tilde{Z}_{n,N}} |\det M|^{2\alpha} e^{-N \text{Tr } V(M)} dM \quad (6.3.18)$$

with $\alpha > -1/2$.

The extra factor $|\det M|^{2\alpha}$ is referred to as a spectral singularity and the matrix model (6.3.18) is relevant in quantum chromodynamics, where it is referred to as a chiral ensemble, see Chapter 32 of this handbook. The spectral singularity does not change the global density ρ_V of eigenvalues, but it does have an influence on the local eigenvalue statistics at the origin. Assuming that $c = \rho_V(0) > 0$, one now finds

$$\lim_{n \rightarrow \infty} \frac{1}{cn} K_{n,N} \left(\frac{x}{cn}, \frac{y}{cn} \right) = \widehat{K}^{\text{Bes}, \alpha}(x, y), \quad x, y > 0, \quad (6.3.19)$$

instead of (6.3.12).

The universality of the Bessel kernels (6.2.5) and (6.3.17) was discussed by Akemann et al. [Ake97b] for integer values of α . It was extended to non-integer α in [Kan98]. The model (6.3.18) was analyzed in [Kui03] with the Riemann-Hilbert method.

See [Kle00] for the spectral universality in orthogonal ensembles.

Weight with jump discontinuity More recently [Fou11], a new class of limiting kernels was identified for unitary ensembles of the form

$$\frac{1}{\tilde{Z}_n} e^{-\text{Tr } V(M)} dM$$

defined on Hermitian matrices with eigenvalues in $[-1, 1]$ in cases where the weight function $w(x) = e^{-V(x)}$, $x \in [-1, 1]$, has a jump discontinuity at $x = 0$. The limiting kernels in [Fou11] are constructed out of confluent hypergeometric functions, see also [Its08].

6.4 Riemann-Hilbert method

A variety of methods have been developed to prove the above universality results in varying degrees of generality. One of these methods will be described here, namely the steepest descent analysis for the Riemann-Hilbert problem (RH problem). This is a powerful method to obtain strong and uniform asymptotics for orthogonal polynomials. One of the outcomes is the limiting behavior of the eigenvalue correlation kernels. However, the method gives much more. It is also able to give asymptotics of the recurrence coefficients, Hankel determinants and other notions associated with orthogonal polynomials. In this section we closely follow the paper [Dei99c] of Deift et al, see also [Dei99a].

6.4.1 Statement of the RH problem

A Riemann-Hilbert problem is a jump problem for a piecewise analytic function in the complex plane. The RH problem for orthogonal polynomials asks for a 2×2 matrix valued function Y satisfying

- (1) $Y : \mathbb{C} \setminus \mathbb{R} \rightarrow \mathbb{C}^{2 \times 2}$ is analytic,
- (2) Y has boundary values on the real line, denoted by $Y_{\pm}(x)$, where $Y_+(x)$ ($Y_-(x)$) denotes the limit of $Y(z)$ as $z \rightarrow x \in \mathbb{R}$ with $\text{Im } z > 0$ ($\text{Im } z < 0$), and

$$Y_+(x) = Y_-(x) \begin{pmatrix} 1 & e^{-NV(x)} \\ 0 & 1 \end{pmatrix},$$

- (3) $Y(z) = (I + O(1/z)) \begin{pmatrix} z^n & 0 \\ 0 & z^{-n} \end{pmatrix}$ as $z \rightarrow \infty$.

The unique solution is given in terms of the orthogonal polynomials (6.3.4) by

$$Y(z) = \begin{pmatrix} P_{n,N}(z) & \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{P_{n,N}(x)e^{-NV(x)}}{x-z} dx \\ -2\pi i \gamma_{n-1,N}^2 P_{n-1,N}(z) & -\gamma_{n-1,N}^2 \int_{-\infty}^{\infty} \frac{P_{n-1,N}(x)e^{-NV(x)}}{x-z} dx \end{pmatrix}. \quad (6.4.1)$$

The RH problem for orthogonal polynomials and its solution (6.4.1) are due to Fokas, Its, and Kitaev [Fok92].

By the Christoffel-Darboux formula for orthogonal polynomials, we have that the orthogonal polynomial kernel (6.3.5) is equal to

$$K_{n,N}(x, y) = \sqrt{e^{-NV(x)}} \sqrt{e^{-NV(y)}} \gamma_{n-1,N}^2 \frac{P_{n,N}(x)P_{n-1,N}(y) - P_{n-1,N}(x)P_{n,N}(y)}{x-y} \quad (6.4.2)$$

which in view of (6.4.1) and the fact that $\det Y(z) \equiv 1$ can be rewritten as

$$K_{n,N}(x, y) = \frac{1}{2\pi i(x-y)} \sqrt{e^{-NV(x)}} \sqrt{e^{-NV(y)}} \begin{pmatrix} 0 & 1 \end{pmatrix} Y_+(y)^{-1} Y_+(x) \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (6.4.3)$$

for $x, y \in \mathbb{R}$. Other notions related to the orthogonal polynomials are also contained in the solution of the Riemann-Hilbert problem. The monic polynomials satisfy a three-term recurrence relation

$$xP_{n,N}(x) = P_{n+1,N}(x) + b_{n,N}P_{n,N}(x) + a_{n,N}P_{n-1,N}(x)$$

with recurrence coefficients $a_{n,N} > 0$ and $b_{n,N} \in \mathbb{R}$. The recurrence coefficients can be found from the solution of the RH problem by expanding Y around ∞ :

$$Y(z) = \left(I + \frac{1}{z}Y_1 + \frac{1}{z^2}Y_2 + \cdots \right) \begin{pmatrix} z^n & 0 \\ 0 & z^{-n} \end{pmatrix}$$

as $z \rightarrow \infty$. The 2×2 matrices Y_1 and Y_2 do not depend on z , but do depend on n and N . Then

$$a_{n,N} = (Y_1)_{12}(Y_1)_{21}, \quad b_{n,N} = \frac{(Y_2)_{12}}{(Y_1)_{12}} - (Y_1)_{22}. \quad (6.4.4)$$

6.4.2 Outline of the steepest descent analysis

The steepest descent analysis of RH problems is due to Deift and Zhou [Dei93]. It produces a number of explicit and invertible transformations leading to a RH problem for a new matrix-valued function R with identity asymptotics at infinity. Also R depends on n and N , and as $n, N \rightarrow \infty$ with $n/N \rightarrow 1$, the jump matrices tend to the identity matrix, in regular cases typically at a rate of $O(1/n)$. Then it can be shown that $R(z)$ tends to the identity matrix as $n, N \rightarrow \infty$ with $n/N \rightarrow 1$ and also at a rate of $O(1/n)$ in regular cases.

Following the transformations in the steepest descent analysis we can then find asymptotic formulas for Y and in particular for the orthogonal polynomial $P_{n,N}$, the recurrence coefficients and the correlation kernel.

With more work one may be able to obtain more precise asymptotic information on R . For example, if $n = N$ and if we are in the one-cut regular case, then there is an asymptotic expansion for $R(z)$:

$$R(z) \sim I + \frac{R^{(1)}(z)}{n} + \frac{R^{(2)}(z)}{n^2} + \cdots$$

with explicitly computable matrices $R^{(j)}(z)$. This in turn leads to asymptotic expansions for the orthogonal polynomials as well. We will not go into this aspect here.

Here we present the typical steps in the Deift-Zhou steepest descent analysis. We focus on the one-cut case, that is on the situation where V is real analytic and the equilibrium measure μ_V is supported on one interval $[a, b]$. We also assume that we are in a regular case, which means that

$$\rho_V(x) = \frac{d\mu_V(x)}{dx} = \frac{1}{\pi} h(x) \sqrt{(b-x)(x-a)}, \quad x \in [a, b],$$

with $h(x) > 0$ for $x \in [a, b]$, and strict inequality holds in the variational condition (6.4.6) below. Generically these regularity conditions are satisfied, see [Kui00]. The singular cases lead to different universality classes, and this will be commented on in section 6.5.

For convenience we also take $N = n$.

6.4.3 First transformation: normalization at infinity

The equilibrium measure μ_V is used in the first transformation of the RH problem. The equilibrium measure satisfies for some constant ℓ

$$2 \int \log \frac{1}{|x-y|} d\mu_V(y) + V(x) = \ell, \quad x \in [a, b], \quad (6.4.5)$$

$$2 \int \log \frac{1}{|x-y|} d\mu_V(y) + V(x) \geq \ell, \quad x \in \mathbb{R} \setminus [a, b]. \quad (6.4.6)$$

These are the Euler-Lagrange variational conditions associated with the minimization of the weighted energy (6.3.8). Since (6.3.8) is a strictly convex functional on finite energy probability measures, the conditions (6.4.5)–(6.4.6) characterize the minimizer μ_V .

The equilibrium measure μ_V leads to the g -function

$$g(z) = \int \log(z-x) d\mu_V(x), \quad z \in \mathbb{C} \setminus (-\infty, b], \quad (6.4.7)$$

which is used in the first transformation $Y \mapsto T$. We put

$$T(z) = \begin{pmatrix} e^{-n\ell/2} & 0 \\ 0 & e^{n\ell/2} \end{pmatrix} Y(z) \begin{pmatrix} e^{-n(g(z)-\ell/2)} & 0 \\ 0 & e^{n(g(z)-\ell/2)} \end{pmatrix}. \quad (6.4.8)$$

The jumps in the RH problem for T are conveniently stated in terms of the functions

$$\phi(z) = \int_b^z h(s)((s-b)(s-a))^{1/2} ds \quad (6.4.9)$$

$$\tilde{\phi}(z) = \int_a^z h(s)((s-b)(s-a))^{1/2} ds, \quad (6.4.10)$$

where we use the fact that V is real analytic and therefore h has an analytic continuation to a neighborhood of the real line. Then T satisfies the RH problem




Figure 6.1: Contour Σ_S and jump matrices for the RH problem for S . This figure is reproduced from [Kui09].

- (1) T is analytic in $\mathbb{C} \setminus \mathbb{R}$.
- (2) On $\mathbb{R}$ we have the jump $T_+ = T_- J_T$ where

$$J_T(x) = \begin{cases} \begin{pmatrix} e^{2n\phi_+(x)} & 1 \\ 0 & e^{2n\phi_-(x)} \end{pmatrix} & x \in (a, b), \\ \begin{pmatrix} 1 & e^{-2n\phi(x)} \\ 0 & 1 \end{pmatrix} & x > b, \\ \begin{pmatrix} 1 & e^{-2n\tilde{\phi}(x)} \\ 0 & 1 \end{pmatrix} & x < a. \end{cases} \quad (6.4.11)$$

- (3) $T(z) = I + O(1/z)$ as $z \rightarrow \infty$.

The RH problem is now normalized at infinity. The jump matrices on $(-\infty, a)$ and (b, ∞) tend to the identity matrix as $n \rightarrow \infty$, since $\phi(x) > 0$ for $x > b$ and $\tilde{\phi}(x) > 0$ for $x < a$, which is due to the assumption that strict inequality holds in (6.4.6). We have to deal with the jump on (a, b) . The diagonal entries in the jump matrix on (a, b) are highly oscillatory. The goal of the next transformation is to turn these highly oscillatory entries into exponentially decaying ones.

6.4.4 Second transformation: opening of lenses

We open up a lens around the interval $[a, b]$ as in Figure 6.1 and define

$$S = \begin{cases} T \begin{pmatrix} 1 & 0 \\ -e^{2n\phi} & 0 \end{pmatrix} & \text{in the upper part of the lens,} \\ T \begin{pmatrix} 1 & 0 \\ e^{2n\phi} & 0 \end{pmatrix} & \text{in the lower part of the lens} \\ T & \text{elsewhere.} \end{cases} \quad (6.4.12)$$

Then S is defined and analytic in $\mathbb{C} \setminus \Sigma_S$ where Σ_S is the contour consisting of the real line, and the upper and lower lips of the lens, with orientation as indicated by the arrows in Figure 6.1. The orientation induces a $+$ -side and a $-$ -side on each part of Σ_S , where the $+$ -side is on the left as one traverses the contour according to its orientation and the $-$ -side is on the right. We use $S_{\pm}$ to denote the limiting values of S on Σ_S when approaching Σ_S from the $\pm$ -side. This convention about $\pm$ -limits, depending on the orientation of the contour is usual in Riemann-Hilbert problems and will also be used later on.

Then S satisfies the following RH problem:

- (1) S is analytic in $\mathbb{C} \setminus (\mathbb{R} \cup \Sigma_S)$.
- (2) On Σ_S we have the jump $S_+ = S_- J_S$ where

$$J_S(x) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } x \in (a, b), \\ \begin{pmatrix} 1 & 0 \\ e^{2n\phi(x)} & 1 \end{pmatrix} & \text{on the lips of the lens,} \\ J_T(x) & \text{for } x < a \text{ or } x > b. \end{cases} \quad (6.4.13)$$

- (3) $S(z) = I + O(1/z)$ as $z \rightarrow \infty$.

If the lens is taken sufficiently small, we can guarantee that $\operatorname{Re} \phi < 0$ on the lips of the lens. Then the jump matrices for S tend to the identity matrix as $n \rightarrow \infty$ on the lips of the lens and on the unbounded intervals $(-\infty, a)$ and (b, ∞) .

6.4.5 Outside parametrix

The next step is to build an approximation to S , valid for large n , the so-called parametrix. The parametrix consists of two parts, an outside or global parametrix that will model S away from the endpoints and local parametrices that are good approximation to S in a neighborhood of the endpoints.

The outside parametrix M satisfies

- (1) M is analytic in $\mathbb{C} \setminus [a, b]$,
- (2) M has the jump

$$M_+(x) = M_-(x) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{for } x \in (a, b).$$

- (3) $M(z) = I + O(1/z)$ as $z \rightarrow \infty$.

To have uniqueness of a solution we also impose




Figure 6.2: Neighborhood $U_\delta(b)$ of b and contours and jump matrices for the RH problem for P

- (4) M has at most fourth-root singularities at the endpoints a and b .

The solution to this RH problem is given by

$$M(z) = \begin{pmatrix} \frac{\beta(z) + \beta^{-1}(z)}{2} & \frac{\beta(z) - \beta^{-1}(z)}{2i} \\ -\frac{\beta(z) - \beta^{-1}(z)}{2i} & \frac{\beta(z) + \beta^{-1}(z)}{2} \end{pmatrix}, \quad z \in \mathbb{C} \setminus [a, b], \quad (6.4.14)$$

where

$$\beta(z) = \left(\frac{z-b}{z-a} \right)^{1/4}. \quad (6.4.15)$$

6.4.6 Local parametrix

The local parametrix is constructed in neighborhoods $U_\delta(a) = \{z \mid |z-a| < \delta\}$ and $U_\delta(b) = \{z \mid |z-b| < \delta\}$ of the endpoints a and b , where $\delta > 0$ is small, but fixed.

In $U_\delta(a) \cup U_\delta(b)$ we want to have a 2×2 matrix valued function P satisfying the following (see also Figure 6.2 for the jumps in the neighborhood of b):

- (1) P is defined and analytic in $(U_\delta(a) \cup U_\delta(b)) \setminus \Sigma_S$ and has a continuous extension to $(\overline{U_\delta(a)} \cup \overline{U_\delta(b)}) \setminus \Sigma_S$.
- (2) On $\Sigma_S \cap (U_\delta(a) \cup U_\delta(b))$ there is the jump

$$P_+ = P_- J_S$$




Figure 6.3: Contour Σ_A and jump matrices for the Airy Riemann-Hilbert problem.

where J_S is the jump matrix (6.4.13) in the RH problem for S .

- (3) P agrees with the global parametrix M on the boundaries of $U_\delta(a)$ and $U_\delta(b)$ in the sense that

$$P(z) = \left(I + O\left(\frac{1}{n}\right) \right) M(z) \quad (6.4.16)$$

as $n \rightarrow \infty$ uniformly for $z \in \partial U_\delta(a) \cup \partial U_\delta(b)$.

- (4) $P(z)$ remains bounded as $z \rightarrow a$ or $z \rightarrow b$.

The solution of the RH problem for P is constructed out of the Airy function

$$y_0(z) = \text{Ai}(z),$$

and its rotated versions

$$y_1(z) = \omega \text{Ai}(\omega z), \quad y_2(z) = \omega^2 \text{Ai}(\omega^2 z), \quad \omega = e^{2\pi i/3}.$$

These three solutions of the Airy differential equation $y''(z) = zy(z)$ are connected by the identity

$$y_0(z) + y_1(z) + y_2(z) = 0.$$

They are used in the solution of the following model RH problem posed on the contour Σ_A shown in Figure 6.3.

- (1) $A : \mathbb{C} \setminus \Sigma_A \rightarrow \mathbb{C}^{2 \times 2}$ is analytic.

(2) A has jump $A_+ = A_- J_A$ on Σ_A with jump matrix J_A given by

$$J_A(z) = \begin{cases} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } \arg z = 0, \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } \arg z = \pm 2\pi i/3, \\ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } \arg z = \pi. \end{cases} \quad (6.4.17)$$

(3) As $z \rightarrow \infty$, we have

$$A(z) = \begin{pmatrix} z^{-1/4} & 0 \\ 0 & z^{1/4} \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix} (I + O(z^{-3/2})) \begin{pmatrix} e^{-\frac{2}{3}z^{3/2}} & 0 \\ 0 & e^{\frac{2}{3}z^{3/2}} \end{pmatrix}. \quad (6.4.18)$$

(4) $A(z)$ remains bounded as $z \rightarrow 0$.

The rather complicated asymptotics in (6.4.18) corresponds to the asymptotic formulas

$$\begin{aligned} \text{Ai}(z) &= \frac{1}{2\sqrt{\pi}z^{1/4}} e^{-\frac{2}{3}z^{3/2}} \left(1 + \mathcal{O}(z^{-3/2})\right), \\ \text{Ai}'(z) &= -\frac{z^{1/4}}{2\sqrt{\pi}} e^{-\frac{2}{3}z^{3/2}} \left(1 + \mathcal{O}(z^{-3/2})\right), \end{aligned} \quad (6.4.19)$$

as $z \rightarrow 0$ with $-\pi < \arg z < \pi$, that are known for the Airy function and its derivative.


The solution of the Airy Riemann-Hilbert problem is as follows

$$A(z) = \sqrt{2\pi} \times \begin{cases} \begin{pmatrix} y_0(z) & -y_2(z) \\ -iy'_0(z) & iy'_2(z) \end{pmatrix}, & 0 < \arg z < 2\pi/3, \\ \begin{pmatrix} -y_1(z) & -y_2(z) \\ -iy'_1(z) & iy'_2(z) \end{pmatrix}, & 2\pi/3 < \arg z < \pi, \\ \begin{pmatrix} -y_2(z) & y_1(z) \\ iy'_2(z) & -iy'_1(z) \end{pmatrix}, & -\pi < \arg z < -2\pi/3, \\ \begin{pmatrix} y_0(z) & y_1(z) \\ -iy'_0(z) & -iy'_1(z) \end{pmatrix}, & -2\pi/3 < \arg z < 0. \end{cases} \quad (6.4.20)$$

The constants $\sqrt{2\pi}$ and $\pm i$ are such that $\det A(z) \equiv 1$ for $z \in \mathbb{C} \setminus \Sigma_A$.

To construct the local parametrix P in the neighborhood $U_\delta(b)$ of b we also need the function

$$f(z) = \left[\frac{3}{2} \phi(z) \right]^{2/3}$$

Figure 6.4: Contour Σ_R and jump matrices for the RH problem for R .

which is a conformal map from $U_\delta(b)$ (provided δ is small enough) onto a neighborhood of the origin. For this it is important that the density ρ_V vanishes as a square root at b . We may assume that the lens around (a, b) is opened in such a way that the parts of the lips of the lens within $U_\delta(b)$ are mapped by f into the rays $\arg z = \pm 2\pi/3$.

Then the local parametrix P is given in $U_\delta(b)$ by

$$P(z) = E_n(z)A(n^{2/3}f(z)) \begin{pmatrix} e^{n\phi(z)} & 0 \\ 0 & e^{-n\phi(z)} \end{pmatrix}, \quad z \in U_\delta(b) \setminus \Sigma_S, \quad (6.4.21)$$

where the prefactor $E_n(z)$ is given explicitly by

$$E_n(z) = -\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix} \begin{pmatrix} n^{1/6}(z-a)^{1/4} & 0 \\ 0 & n^{-1/6}(z-a)^{-1/4} \end{pmatrix} \times \begin{pmatrix} (f(z)/(z-b))^{1/4} & 0 \\ 0 & (f(z)/(z-b))^{-1/4} \end{pmatrix}. \quad (6.4.22)$$

Then E_n is analytic in $U_\delta(b)$ and it does not change the jump conditions. It is needed in order to satisfy the matching condition (6.4.16) on the boundary $|z-b|=\delta$ of $U_\delta(b)$.

A similar construction gives the local parametrix P in the neighborhood $U_\delta(a)$ of a .

6.4.7 Final transformation

In the final transformation we put

$$R(z) = \begin{cases} S(z)M(z)^{-1}, & z \in \mathbb{C} \setminus (\Sigma_S \cup \overline{U_\delta(a)} \cup \overline{U_\delta(b)}), \\ S(z)P(z)^{-1}, & z \in (U_\delta(a) \cup U_\delta(b)) \setminus \Sigma_S. \end{cases} \quad (6.4.23)$$

Then R has an analytic continuation to $\mathbb{C} \setminus \Sigma_R$ where Σ_R is shown in Figure 6.4.

R solves the following RH problem.

- (1) R is analytic in $\mathbb{C} \setminus \Sigma_R$.
- (2) R satisfies the jump conditions $R_+ = R_- J_R$ where J_R are the matrices given in Figure 6.4.
- (3) $R(z) = I + O(z^{-1})$ as $z \rightarrow \infty$.

The jump matrices tend to the identity matrix as $n \rightarrow \infty$. Indeed, the jump matrix J_R on the boundaries of the disks satisfies

$$J_R(z) = P(z)M(z)^{-1} = I + O(n^{-1})$$

as $n \rightarrow \infty$, because of the matching condition (6.4.16). On the remaining parts of Σ_R we have that $J_R = I + O(e^{-cn})$ as $n \rightarrow \infty$ for some constant $c > 0$.

Technical estimates on solutions of RH problems, see [Dei99a], now guarantee that in this case

$$R(z) = I + O(1/n) \quad \text{as } n \rightarrow \infty, \quad (6.4.24)$$

uniformly for $z \in \mathbb{C} \setminus \Sigma_R$.

6.4.8 Proof of bulk universality

Now we turn our attention again to the correlation kernel (6.4.3) which, since $n = N$, we denote by K_n instead of $K_{n,N}$. We follow what happens with K_n under the transformations $Y \mapsto T \mapsto S$. We assume that $x, y \in (a, b)$.

From the definition (6.4.8) we obtain

$$K_n(x, y) = \frac{\sqrt{e^{-nV(x)}}\sqrt{e^{-nV(y)}}}{2\pi i(x-y)} \begin{pmatrix} 0 & e^{n(g_+(y)+\ell/2)} \end{pmatrix} T_+^{-1}(y)T_+(x) \begin{pmatrix} e^{n(g_+(x)+\ell/2)} \\ 0 \end{pmatrix}$$

which by (non-trivial) properties of (6.4.7) and (6.4.9), based on the variational condition (6.4.5), can be rewritten as

$$K_n(x, y) = \frac{1}{2\pi i(x-y)} \begin{pmatrix} 0 & e^{-n\phi_+(y)} \end{pmatrix} T_+^{-1}(y)T_+(x) \begin{pmatrix} e^{-n\phi_+(x)} \\ 0 \end{pmatrix}. \quad (6.4.25)$$

Using the transformation (6.4.12) in the upper part of the lens, we see that (6.4.25) leads to

$$K_n(x, y) = \frac{1}{2\pi i(x-y)} \begin{pmatrix} -e^{n\phi_+(y)} & e^{-n\phi_+(y)} \end{pmatrix} S_+^{-1}(y)S_+(x) \begin{pmatrix} e^{-n\phi_+(x)} \\ e^{n\phi_+(x)} \end{pmatrix} \quad (6.4.26)$$

for $x, y \in (a, b)$. This is the basic formula for K_n in terms of S .

If $x, y \in (a + \delta, b - \delta)$, then by (6.4.23)

$$S_+^{-1}(y)S_+(x) = M_+^{-1}(y)R(y)^{-1}R(x)M_+(x).$$

The uniform estimate (6.4.24) on R can then be used to show that

$$S_+^{-1}(y)S_+(x) = I + O(x - y). \quad (6.4.27)$$

with an O -term that is uniform for $x, y \in (a + \delta, b - \delta)$. Using (6.4.27) in (6.4.26) we arrive at

$$K_n(x, y) = \frac{\sin(in(\phi_+(y) - \phi_+(x)))}{\pi(x - y)} + O(1). \quad (6.4.28)$$

Take $x^* \in (a, b)$ fixed and let $c = \rho_V(x^*) > 0$. We may assume that $\delta > 0$ has been chosen so small that $x^* \in (a + \delta, b - \delta)$. Replace x and y in (6.4.28) by $x^* + \frac{x}{cn}$ and $x^* + \frac{y}{cn}$ respectively. Then after dividing through by cn , we have for fixed x and y ,

$$\begin{aligned} \frac{1}{cn} K_n \left(x^* + \frac{x}{cn}, x^* + \frac{y}{cn} \right) \\ = \frac{\sin \left(in \left(\phi_+ \left(x^* + \frac{y}{cn} \right) - \phi_+ \left(x^* + \frac{x}{cn} \right) \right) \right)}{\pi(x - y)} + O \left(\frac{1}{n} \right) \end{aligned} \quad (6.4.29)$$

as $n \rightarrow \infty$, and the O -term is uniform for x and y in compact subsets of $\mathbb{R}$. Since $\phi'_+(s) = \pi i \rho_V(s)$ for $s \in (a, b)$, and $c = \rho_V(x^*)$, we have that

$$in \left(\phi_+ \left(x^* + \frac{y}{cn} \right) - \phi_+ \left(x^* + \frac{x}{cn} \right) \right) = \pi(x - y) + O \left(\frac{x - y}{n} \right)$$

and therefore the rescaled kernel (6.4.29) does indeed tend to the sine kernel (6.2.3) as $n \rightarrow \infty$. This proves the bulk universality.

Remark. Letting $y \rightarrow x$ in (6.4.28) and dividing by n we also obtain

$$\begin{aligned} \frac{1}{n} K_n(x, x) &= \frac{1}{\pi i} \phi'_+(x) + O \left(\frac{1}{n} \right) \\ &= \frac{1}{\pi} h(x) \sqrt{(b - x)(x - a)} + O \left(\frac{1}{n} \right) \quad (\text{by (6.4.9)}) \\ &= \rho_V(x) + O \left(\frac{1}{n} \right) \quad (\text{by (6.3.9)}) \end{aligned}$$

as $n \rightarrow \infty$, uniformly for x in compact subsets of (a, b) , which proves that ρ_V is the limiting means eigenvalue density.

6.4.9 Proof of edge universality

We will not give the proof of the edge universality in detail. The proof starts from the representation (6.4.26) of the correlation kernel in terms of S . In the neighborhood of a and b we have by (6.4.23) that $S = RP$, where P is the local

parametrix that is constructed out of Airy functions, as it involves the solution of the Airy Riemann-Hilbert problem.

From (6.4.20) it can be checked that the Airy kernel (6.2.4) is given in terms of the solution $A(z)$ by

$$K^{\text{Ai}}(x, y) = \frac{1}{2\pi i(x - y)} \times \begin{cases} \begin{pmatrix} 0 & 1 \end{pmatrix} A_+^{-1}(y) A_+(x) \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \text{if both } x, y > 0, \\ \begin{pmatrix} -1 & 1 \end{pmatrix} A_+^{-1}(y) A_+(x) \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \text{if } x > 0 \text{ and } y < 0, \\ \begin{pmatrix} 0 & 1 \end{pmatrix} A_+^{-1}(y) A_+(x) \begin{pmatrix} 1 \\ 1 \end{pmatrix} & \text{if } x < 0 \text{ and } y > 0, \\ \begin{pmatrix} -1 & 1 \end{pmatrix} A_+^{-1}(y) A_+(x) \begin{pmatrix} 1 \\ 1 \end{pmatrix} & \text{if both } x, y < 0, \end{cases}$$

and this is exactly what comes out of the calculations for the scaling limit of the eigenvalue correlation kernels K_n near the edge point.

6.5 Non-standard universality classes

The standard universality classes (sine, Airy and Bessel) describe the local eigenvalue statistics around regular points.

In the unitary ensemble (6.3.1) there are three types of singular eigenvalue behavior. They all depend on the behavior of the global eigenvalue density ρ_V . The three types of singular behavior are:

- The density ρ_V vanishes at an interior point of the support.
- The density ρ_V vanishes to higher order at an edge point of the support (higher than square root).
- Equality holds in the variational inequality (6.4.6) at a point outside $\text{supp}(\mu_V)$.

In each of these cases there exists a family of limiting correlation kernels that arise in a double scaling limit. In (6.3.1) and (6.3.3) one lets $n, N \rightarrow \infty$ with $n/N \rightarrow 1$ at a critical rate so that for some exponent γ , the limit

$$\lim_{n \rightarrow \infty} n^\gamma \left(\frac{n}{N} - 1 \right) \quad (6.5.1)$$

exists. The family of limiting kernels is parametrized by the value of the limit (6.5.1).

Most rigorous results that have been obtained in this direction are based on the RH problem, and use an extension of the steepest descent analysis that was discussed in the previous section. Non-standard universality classes are also discussed in Chapters 12 and 13.

6.5.1 Interior singular point

An interior singular point is a point x^* where ρ_V vanishes. Varying the parameters in V may then either lead to a gap in the support around x^* , or to the closing of the gap. The local scaling limits at x^* depend on the order of vanishing at x^* , that is, on the positive integer k so that

$$\rho_V(x) = c(x - x^*)^{2k}(1 + o(1)) \quad \text{as } x \rightarrow x^*$$

with $c > 0$.

The case of quadratic vanishing (i.e., $k = 1$) was considered in [Ble03] for the critical quartic potential

$$V(x) = \frac{1}{4}x^4 - x^2$$

and in [Cla06, Shc08] for more general V . For $k = 1$ one takes $\gamma = 2/3$ in (6.5.1). The limiting kernels are parametrized by a parameter s which is proportional to the limit (6.5.1)

$$s = c \lim_{n \rightarrow \infty} n^{2/3} \left(\frac{n}{N} - 1 \right),$$

where the proportionality constant $c > 0$ is (for the case $\text{supp}(\mu_V) = [a, b]$)

$$c = \frac{2}{(\pi \rho_V''(x^*))^{1/3} \sqrt{(b - x^*)(x^* - a)}}.$$

The s -dependence is then governed by the Hastings-McLeod solution $q(s)$ of the Painlevé II equation

$$q'' = sq + 2q^3. \quad (6.5.2)$$

The limiting kernels are therefore called Painlevé II kernels and we denote them by $K^{\text{PII}}(x, y; s)$, even though $q(s)$ itself does not appear in the formulas for the kernels. What does appear is a solution of the Lax pair equations

$$\begin{aligned} \frac{\partial}{\partial x} \begin{pmatrix} \Phi_1(x; s) \\ \Phi_2(x; s) \end{pmatrix} &= \begin{pmatrix} -4ix^2 - i(s + 2q^2) & 4xq + 2ir \\ 4xq - 2ir & 4ix^2 + i(s + 2q^2) \end{pmatrix} \begin{pmatrix} \Phi_1(x; s) \\ \Phi_2(x; s) \end{pmatrix} \\ \frac{\partial}{\partial s} \begin{pmatrix} \Phi_1(x; s) \\ \Phi_2(x; s) \end{pmatrix} &= \begin{pmatrix} -ix & q \\ q & ix \end{pmatrix} \begin{pmatrix} \Phi_1(x; s) \\ \Phi_2(x; s) \end{pmatrix} \end{aligned} \quad (6.5.3)$$

where $q = q(s)$ is the Hastings-McLeod solution of (6.5.2) and $r = r(s) = q'(s)$.

There is a specific solution of (6.5.3) so that the family of limiting kernels takes the form

$$K^{\text{PII}}(x, y; s) = \frac{-\Phi_1(x; s)\Phi_2(y; s) + \Phi_1(y; s)\Phi_2(x; s)}{2\pi i(x - y)}. \quad (6.5.4)$$

For $k \geq 2$, one takes $\gamma = 2k/(2k + 1)$ in (6.5.1), and the limiting kernels can be described in a similar way by a special solution of the k th member of

the Painlevé II hierarchy. The functions in the kernels itself are solutions of the associated Lax pair equations.

The connection with Painlevé II also holds for the unitary matrix model with a spectral singularity (6.3.18)

$$\frac{1}{\tilde{Z}_{n,N}} |\det M|^{2\alpha} e^{-N \operatorname{Tr} V(M)} dM, \quad \alpha > -1/2.$$

In the multicritical case where ρ_V vanishes quadratically at $x = 0$, the limiting kernels are associated with a special solution of the general form of the Painlevé II equation

$$q'' = sq + 2q^3 - \alpha$$

with parameter α , see [Ake98, Cla08].

6.5.2 Singular edge point

A singular edge point is a point x^* where ρ_V vanishes to higher order than square root. The local scaling limits depend again on the order of vanishing. If x^* is a right-edge point of an interval in the support, then there is an even integer k such that

$$\rho_V(x) = c(x^* - x)^{k+1/2}(1 + o(1)) \quad \text{as } x \nearrow x^* \quad (6.5.5)$$

with $c > 0$. Here one takes $\gamma = (2k + 2)/(2k + 3)$ in (6.5.1). The limiting kernels are described by Lax pair solutions associated with a special solution of the k th member of the Painlevé I hierarchy. For $k = 2$ this is worked out in detail in [Cla07]. For general k , and assuming x^* is the right-most point in the support, the largest eigenvalue distributions are studied in [Cla09]. These generalizations of the Tracy-Widom distribution are expressed in terms of members of a Painlevé II hierarchy.

The case $k = 1$ in (6.5.5) cannot occur in the unitary matrix model (6.3.1), although it frequently appears in the physics literature, see e.g. [DiF95], where, for example, it is associated with the potential

$$V(x) = -\frac{1}{48}x^4 + \frac{1}{2}x^2.$$

This potential does not satisfy (6.3.2) and the model (6.3.1) is not well-defined as a probability measure on Hermitian matrices, although it can be studied in a formal sense, see e.g. in Chapter 16.

In [Fok92] and [Dui06] the polynomials that are orthogonal on certain contours in the complex plane with a weight $e^{-NV(x)}$, where $V(x) = t\frac{x^4}{4} + \frac{x^2}{2}$ and $t \approx -\frac{1}{12}$, are studied and the connection with special solutions of the Painlevé I equation

$$q'' = 6q^2 + s$$

is rigorously established.

6.5.3 Exterior singular point

An exterior point x^* is singular if there is equality in the variational inequality (6.4.6). At such a point a new interval in the support may arise when perturbing the potential V . The limiting kernels at such a point depend on the order of vanishing of

$$2 \int \log \frac{1}{|x-y|} d\mu_V(y) + V(x) - \ell$$

at $x = x^*$.

In this situation the appropriate scaling is so that

$$\lim_{n \rightarrow \infty} \frac{n}{\log n} \left(\frac{n}{N} - 1 \right) \quad (6.5.6)$$

exists, instead of (6.5.1). This special kind of scaling was discussed in [Ake97a, Eyn06]. Maybe surprisingly, there is no connection with Painlevé equations in this case. In the simplest case of quadratic vanishing at $x = x^*$ the possible limiting kernels are finite size GUE kernels and certain interpolants, see [Ber09, Cla08, Mo08].

6.5.4 Pearcey kernels

The Painlevé II kernels (6.5.4) are the canonical kernels that arise at the closing of a gap in unitary ensembles.

Brézin and Hikami [Bre98] were the first to identify a second one parameter family of kernels that may arise at the closing of a gap. This is the family of Pearcey kernels

$$K^{\text{Pear}}(x, y; s) = \frac{p(x)q''(y) - p'(x)q'(y) + p''(x)q(y) - sp(x)q(y)}{x - y} \quad (6.5.7)$$

with $s \in \mathbb{R}$, where p and q are solutions of the Pearcey differential equations $p'''(x) = xp(x) - sp'(x)$ and $q'''(y) = yq(y) + sq'(y)$. The kernel is also given by the double integral

$$K^{\text{Pear}}(x, y; s) = \frac{1}{(2\pi i)^2} \int_C \int_{-i\infty}^{i\infty} e^{-\frac{1}{4}\eta^4 + \frac{s}{2}\eta^2 - \eta y + \frac{1}{4}\xi^4 - \frac{s}{2}\xi^2 + \xi x} \frac{d\eta d\xi}{\eta - \xi} \quad (6.5.8)$$

where the contour C consists of the two rays from $\pm\infty e^{i\pi/4}$ to 0 together with the two rays from 0 to $\pm\infty e^{-i\pi/4}$.

The Pearcey kernels appear at the closing of the gap in the Hermitian matrix model with external source

$$\frac{1}{Z_n} e^{-n \text{Tr}(V(M) - AM)} dM \quad (6.5.9)$$

where the external source A is a given Hermitian $n \times n$ matrix. This was proved in [Bre98] for the case where $V(x) = \frac{1}{2}x^2$ and A is a diagonal matrix

$$A = \text{diag}(\underbrace{a, \dots, a}_{n/2 \text{ times}}, \underbrace{-a, \dots, -a}_{n/2 \text{ times}}) \quad (6.5.10)$$

depending on the parameter $a > 0$. For $a > 1$ the eigenvalues of M accumulate on two intervals as $n \rightarrow \infty$ and for $0 < a < 1$ on one interval. The Pearcey kernels (6.5.8) appear in a double scaling limit around the critical value $a = 1$, see also [Ble07] for an analysis of an associated 3×3 matrix valued RH problem. The matrix model with external source (6.5.9) with quadratic potential has an interesting interpretation in terms of non-intersecting Brownian motions [Apt05].

The case of a quartic polynomial potential $V(x) = \frac{1}{4}x^4 - \frac{t}{2}x^2$ was analyzed recently in [Ble11]. Here it was found that the closing of the gap can be either of the Pearcey type or of the Painlevé II type, depending on the value of $t \in \mathbb{R}$.

References

- [Ake97a] G. Akemann, Universal correlators for multi-arc complex matrix models, Nucl.Phys. B 507 (1997) 475–500.
- [Ake97b] G. Akemann, P.H. Damgaard, U. Magnea, and S. Nishigaki, Universality of random matrices in the microscopic limit and the Dirac operator spectrum, Nucl. Phys. B 487 (1997), 721–738.
- [Ake98] G. Akemann, P. H. Damgaard, U. Magnea, S. M. Nishigaki, Multicritical microscopic spectral correlators of hermitian and complex matrices, Nucl.Phys. B 519 (1998), 682–714.
- [And10] G.W. Anderson, A. Guionnet, and O. Zeitouni, *An Introduction to Random Matrices*, Cambridge Univ. Press, Cambridge, 2010.
- [Apt05] A.I. Aptekarev, P.M. Bleher, and A.B.J. Kuijlaars, Large n limit of Gaussian random matrices with external source II, Comm. Math. Phys. 259 (2005), 367–389.
- [Ben97] G. Ben Arous and A. Guionnet, Large deviations for Wigner’s law and Voiculescu’s non-commutative entropy, Probab. Theory Related Fields 108 (1997), 517–542.
- [Ber09] M. Bertola and S.Y. Lee, First colonization of a spectral outpost in random matrix theory, Constr. Approx. 30 (2009), 225–263.

- [Ble11] P. Bleher, S. Delvaux, and A.B.J. Kuijlaars, Random matrix model with external source and a constrained vector equilibrium problem, *Comm. Pure Appl. Math.* 64 (2011), no. 1, 116–160.
- [Ble99] P. Bleher and A. Its, Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model, *Ann. of Math.* 150 (1999), 185–266.
- [Ble03] P. Bleher and A. Its, Double scaling limit in the random matrix model: the Riemann-Hilbert approach, *Comm. Pure Appl. Math.* 56 (2003), 433–516.
- [Ble07] P.M. Bleher and A.B.J. Kuijlaars, Large n limit of Gaussian random matrices with external source III: double scaling limit, *Comm. Math. Phys.* 270 (2007), 481–517.
- [Bow91] M.J. Bowick and E. Brézin, Universal scaling of the tail of the density of eigenvalues in random matrix models, *Phys. Lett. B* 268 (1991), 21–28.
- [Bre98] E. Brézin and S. Hikami, Universal singularity at the closure of a gap in a random matrix theory, *Phys. Rev. E* 57 (1998), 4140–4149.
- [Bre93] E. Brézin and A. Zee, Universality of the correlations between eigenvalues of large random matrices, *Nuclear Physics B* 402 (1993), 613–627.
- [Cla08] T. Claeys, Birth of a cut in unitary random matrix ensembles. *Int. Math. Res. Notices* 2008, Art. ID rnm166, 40 pp.
- [Cla09] T. Claeys, A.R. Its, and I. Krasovsky, Higher order analogues of the Tracy-Widom distribution and the Painlevé II hierarchy, *Comm. Pure Appl. Math.* 63 (2010), 362–412.
- [Cla06] T. Claeys and A.B.J. Kuijlaars, Universality of the double scaling limit in random matrix models, *Comm. Pure Appl. Math.* 59 (2006), 1573–1603.
- [Cla08] T. Claeys, A.B.J. Kuijlaars, and M. Vanlessen, Multi-critical unitary random matrix ensembles and the general Painlevé II equation, *Ann. of Math.* 168 (2008), 601–641.
- [Cla07] T. Claeys and M. Vanlessen, Universality of a double scaling limit near singular edge points in random matrix models, *Comm. Math. Phys.* 273 (2007), 499–532.

- [Dei99a] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach*, Amer. Math. Soc., Providence, RI, 1999.
- [Dei07d] P. Deift, Universality for mathematical and physical systems, in: International Congress of Mathematicians, Vol. I, 125–152, Eur. Math. Soc., Zürich, 2007.
- [Dei07a] P. Deift and D. Gioev, Universality in random matrix theory for orthogonal and symplectic ensembles, Int. Math. Res. Papers 2007, Art. ID rpm004, 116 pp.
- [Dei07b] P. Deift and D. Gioev, Universality at the edge of the spectrum for unitary, orthogonal, and symplectic ensembles of random matrices, Comm. Pure Appl. Math. 60 (2007), 867–910.
- [Dei09] P. Deift and D. Gioev, *Random Matrix Theory: Invariant Ensembles and Universality*, Courant Lecture Notes in Mathematics, 18, Amer. Math. Soc., Providence, RI, 2009.
- [Dei07c] P. Deift, D. Gioev, T. Kriecherbauer, and M. Vanlessen, Universality for orthogonal and symplectic Laguerre-type ensembles, J. Stat. Phys. 129 (2007), 949–1053.
- [Dei98] P. Deift, T. Kriecherbauer, and K.T-R McLaughlin, New results on the equilibrium measure for logarithmic potentials in the presence of an external field, J. Approx. Theory 95 (1998), 388–475.
- [Dei99b] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, Comm. Pure Appl. Math. 52 (1999), 1491–1552.
- [Dei99c] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, Comm. Pure Appl. Math. 52 (1999), 1335–1425.
- [Dei93] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Ann. of Math. 137 (1993), 295–368.
- [DiF95] P. Di Francesco, P. Ginsparg, and J. Zinn-Justin, 2D gravity and random matrices, Phys. Rep. 254 (1995), 133 pp.
- [Dui06] M. Duits and A.B.J. Kuijlaars, Painlevé I asymptotics for orthogonal polynomials with respect to a varying quartic weight, Nonlinearity 19 (2006), 2211–2245.

- [Erd10a] L. Erdős, S. Péché, J. Ramírez, B. Schlein and H-T. Yau, Bulk universality for Wigner matrices, *Comm. Pure Appl. Math.* 63 (2010), 895–925.
- [Erd10b] L. Erdős, J. Ramírez, B. Schlein, T. Tao, V. Vu, and H-T. Yau, Bulk universality for Wigner hermitian matrices with subexponential decay, *Math. Res. Lett.* 17 (2010), 667–674.
- [Erd10c] L. Erdős, Universality of Wigner random matrices: a survey of recent results, preprint arXiv:1004.0861.
- [Eyn06] B. Eynard, Universal distribution of random matrix eigenvalues near the “birth of a cut” transition, *J. Stat. Mech. Theory Exp.* 2006, P07005, 33 pp.
- [Fok92] A.S. Fokas, A.R. Its, and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, *Commun. Math. Phys.* 147 (1992), 395–430.
- [For93] P. J. Forrester, The spectrum edge of random matrix ensembles, *Nucl. Phys. B* 402 (1993), 709–728.
- [For10] P.J. Forrester, *Log-Gases and Random Matrices*, LMS Mongraphs Series Vol. 34, Princeton Univ. Press, Princeton N.J., 2010.
- [Fou11] A. Foulquié Moreno, A. Martínez-Finkelshtein, and V.L. Sousa, Asymptotics of orthogonal polynomials for a weight with a jump on $[-1, 1]$, *Constr. Approx.* 33 (2011), 219–263.
- [Hac95] G. Hackenbroich and H. A. Weidenmüller, Universality of random-matrix results for non-Gaussian ensembles, *Phys. Rev. Lett.* 74 (1995), 4118–4121.
- [Its08] A. Its and I. Krasovsky, Hankel determinant and orthogonal polynomials for the Gaussian weight with a jump, in: “Integrable Systems and Random Matrices” (Baik et al. eds.), *Contemp. Math.* 458, Amer. Math. Soc., Providence RI, 2008, pp. 215–247.
- [Joh98] K. Johansson, On fluctuations of eigenvalues of random Hermitian matrices. *Duke Math. J.* 91 (1998), 151–204.
- [Kan97] E. Kanzieper and V. Freilikher, Universality in invariant random-matrix models: existence near the soft edge, *Phys. Rev. E* 55 (1997), 3712–3715.

- [Kan98] E. Kanzieper and V. Freilikher, Random matrix models with log-singular level confinement: method of fictitious fermions, *Philos. Magazine B* 77 (1998), 1161–1172.
- [Kle00] B. Klein and J. Verbaarschot, Spectral universality of real chiral random matrix ensembles, *Nucl. Phys. B* 588 (2000), 483–507.
- [Kui00] A.B.J. Kuijlaars and K.T-R McLaughlin, Generic behavior of the density of states in random matrix theory and equilibrium problems in the presence of real analytic external fields, *Comm. Pure Appl. Math.* 53 (2000), 736–785.
- [Kui09] A.B.J. Kuijlaars and P. Tibboel, The asymptotic behaviour of recurrence coefficients for orthogonal polynomials with varying exponential weights, *J. Comput. Appl. Math.* 233 (2009), 775–785.
- [Kui02] A.B.J. Kuijlaars and M. Vanlessen, Universality for eigenvalue correlations from the modified Jacobi unitary ensemble, *Int. Math. Res. Notices* 2002 (2002), 1575–1600.
- [Kui03] A.B.J. Kuijlaars and M. Vanlessen, Universality for eigenvalue correlations at the origin of the spectrum, *Comm. Math. Phys.* 243 (2003), 163–191.
- [Lev08] E. Levin and D.S. Lubinsky, Universality limits in the bulk for varying measures, *Adv. Math.* 219 (2008), 743–779.
- [Lev09] E. Levin and D.S. Lubinsky, Universality limits for exponential weights, *Constr. Approx.* 29 (2009), 247–275.
- [Lub08a] D.S. Lubinsky, Universality limits in the bulk for arbitrary measures with compact support, *J. d’Analyse Math.* 106 (2008), 373–394.
- [Lub08b] D.S. Lubinsky, Universality limits at the hard edge of the spectrum for measures with compact support, *Int. Math. Res. Notices* 2008, Art. ID rnn 099, 39 pp.
- [Lub09a] D.S. Lubinsky, A new approach to universality limits involving orthogonal polynomials, *Ann. of Math.* 170 (2009), 915–939.
- [Lub09b] D.S. Lubinsky, Universality limits for random matrices and de Branges spaces of entire functions, *J. Funct. Anal.* 256 (2009), 3688–3729.
- [Meh04] M.L. Mehta, *Random Matrices*, third edition, Elsevier/Academic Press, Amsterdam, 2004.

- [McL08] K. T-R. McLaughlin and P.D. Miller, The $\bar{\partial}$ steepest descent method for orthogonal polynomials on the real line with varying weights, Int. Math. Res. Notices 2008, Art. ID rnn 075, 66 pp.
- [Mo08] M.Y. Mo, The Riemann-Hilbert approach to double scaling limit of random matrix eigenvalues near the “birth of a cut” transition, Int. Math. Res. Notices 2008, Art. ID rnn042, 51 pp.
- [Nag93] T. Nagao and M. Wadati, Eigenvalue distribution of random matrices at the spectrum edge, J. Phys. Soc. Japan 62 (1993), 3845–3856.
- [Pas97] L. Pastur and M. Shcherbina, Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles, J. Statist. Phys. 86 (1997), 109–147.
- [Pas08] L. Pastur, M. Shcherbina, Bulk universality and related properties of Hermitian matrix models, J. Stat. Phys. 130 (2008), 205–250.
- [Saf97] E.B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag, Berlin, 1997.
- [Shc08] M. Shcherbina, Double scaling limit for matrix models with non analytic potentials, J. Math. Phys. 49 (2008) 033501, 34 pp.
- [Shc09a] M. Shcherbina, On universality for orthogonal ensembles of random matrices, Comm. Math. Phys. 285 (2009), 957–974.
- [Shc09b] M. Shcherbina, Edge universality for orthogonal ensembles of random matrices, J. Stat. Phys. 136 (2009), 35–50.
- [Sos00] A. Soshnikov, Determinantal random point fields, Russian Math. Surveys 55 (2000), 923–975.
- [Sto00] A. Stojanovic, Universality in orthogonal and symplectic invariant matrix models with quartic potential, Math. Phys. Anal. Geom. 3 (2000), 339–373.
- [Tao11] T. Tao and V. Vu, Random matrices: Universality of local eigenvalue statistics, Acta Math. 206 (2011), 127–204.
- [Tra94] C. Tracy and H. Widom, Level-spacing distributions and the Airy kernel, Comm. Math. Phys. 159 (1994), 151–174.
- [Tra98] C. Tracy and H. Widom, Correlation functions, cluster functions, and spacing distributions for random matrices, J. Stat. Phys. 92 (1998), 809–835.